

MEDICAID

	2010 QARR		2009 QARR		2008 QARR	
	Affinity	SWA	Affinity	SWA	Affinity	SWA
<b>Access to Care</b>						
<i>Child and Adolescent Access to PCPs</i>						
12-24 months	97%	96%	94%	95%	95%	95%
25 months - 6 years	95%	93%	90%	92%	87%	91%
7-11 years	93%	95%	92%	95%	91%	94%
12-19 years	87%	92%	88%	91%	87%	90%
Total for Child/Adolescents	87%	U/A	90%	U/A	88%	92%
<i>Adult Access to Preventative/Ambulatory Health Services</i>						
Age 20-44	80%	82%	78%	82%	78%	81%
Age 45-64	86%	89%	85%	88%	85%	88%
Age 65 and older	86%	89%	83%	88%	84%	88%
Total for Adults	82%	U/A	81%	U/A	81%	84%
<b>Pediatric Preventive Care</b>						
Childhood Immunization Status (Age 2) - Combo 2: 431331 /with VZV	Rotated-off	N/A	87%	79%	Rotated Off	N/A
Childhood Immunization Status (Age 2) - Combo 3: 4313311 /with 4 Prevna1	Rotated-off	N/A	81%▲	73%	Rotated Off	N/A
Lead Testing (Age 2)	Rotated-off	N/A	91%	90%	Rotated Off	N/A
Well Child Visits 15 month olds (5 or 6 Visits)	83%▲	77%	80%▲	72%	Rotated Off	N/A
Well Child Visits 3-6 years	74%▼	80%	76%▼	79%	Rotated Off	N/A
Adolescent Well Child Visits 12-21 years	48%▼	56%	49%▼	55%	Rotated Off	N/A
<i>Adolescent Screening &amp; Counseling (Ages 10-17yo, (Revised Measure)</i>						
Sexual Health	67%	60%	63%	51%	Rotated Off	N/A
Depression	59%	52%	51%	43%	Rotated Off	N/A
Tobacco Use	69%	64%	68%	54%	Rotated Off	N/A
Substance Use	68%	60%	67%	52%	Rotated Off	N/A
<i>Weight Assessment/Counseling (Ages 3 to 17,</i>						
BMI Percentile	66%	65%	54%	51%	45%	43%
Nutrition Counseling	74%	71%	73%▲	61%	58%	57%
Physical Activity Counseling	60%	58%	56%▲	48%	45%	43%
Annual Dental Visit 2-21 yo	51%	53%	49%	51%	47%▼	48%
Annual Dental Visit 2-18 yo	52%▼	54%	51%▼	52%	48%▼	49%
Annual Dental Visit 19-21 yo	40%▼	43%	39%▼	42%	38%	38%
<b>Adult Preventive Care</b>						
Adult BMI Assessment	72%	70%	60%	55%	45%	45%
<b>Women's Preventive Care</b>						
Chlamydia Screening in Women (Ages 16 to 24)	68%	68%	71%▲	67%	64%▲	59%
Breast Cancer Screening (Ages 42 to 69)	68%	68%	69%	68%	67%	67%
Cervical Cancer Screening (Ages 21 to 64)	68%▼	72%	Rotated Off	N/A	73%	73%
Timeliness of Prenatal Care (Initial Visit)	95%▲	90%	Rotated Off	N/A	92%▲	89%
Frequency of Ongoing Prenatal Care (>81% expected)	88%▲	74%	Rotated Off	N/A	86%▲	73%
Postpartum Care (3 to 8 weeks after Delivery)	74%	73%	Rotated Off	N/A	74%	73%
<b>Chronic Condition Care</b>						
Use of Appropriate Medications for People with Asthma 5-17 yo (Rev '10: 5-11)	91%	92%	91%	92%	92%	92%
Use of Appropriate Medications for People with Asthma 18-56 yo (Rev '10: 12-50)	87%	88%	88%	88%	90%	89%
Use of Appropriate Medications for People with Asthma 5-56 yo (Rev '10: 5-50)	89%	90%	89%	90%	91%	90%
Appropriate Treatment for Asthma: 3-Controller measure (Rev '10: 5-50)	76%	76%	76%	75%	80%▲	77%
<i>Comprehensive Diabetes Care</i>						
HbA1c Test	Rotated-off	N/A	89%	89%	Rotated Off	N/A
HbA1c Poorly Controlled >9	Rotated-off	N/A	25%▲	33%	Rotated Off	N/A
HbA1c Well Controlled <7	Rotated-off	N/A	44%	37%	Rotated Off	N/A
HbA1c Well Controlled <8 (First year measure)	Rotated-off	N/A	59%	55%	N/A	N/A
LDL-C Test	Rotated-off	N/A	86%	87%	Rotated Off	N/A
LDL-C Well Controlled <100	Rotated-off	N/A	46%	44%	Rotated Off	N/A
Eye Exam	Rotated-off	N/A	64%	63%	Rotated Off	N/A

MEDICAID

	2010 QARR		2009 QARR		2008 QARR	
	Affinity	SWA	Affinity	SWA	Affinity	SWA
Monitor Nephropathy	Rotated-off	N/A	82%	83%	Rotated Off	N/A
Blood Pressure <130/80	Rotated-off	N/A	36%	35%	Rotated Off	N/A
Blood Pressure <140/90	Rotated-off	N/A	73%▲	65%	Rotated Off	N/A
Diabetes Composite: Nephro, Eye Exam, HbA1c Testing & LDLc	Rotated-off	N/A	49%	50%	N/A	N/A
Diabetes Composite: Not Poorly Controlled (= <9.0) & LDLc <100mg dL	Rotated-off	N/A	38%	36%	N/A	N/A
<i>Cholesterol Mgmt for Patients with Cardiovascular Conditions</i>						
LDL-C Screening	Rotated-off	N/A	90%	91%	Rotated Off	N/A
LDL-C Control <100	Rotated-off	N/A	52%	51%	Rotated Off	N/A
<i>HIV/AIDS Comprehensive Care</i>						
Engaged in Care (2 Visits)	89%▲	80%	83%▲	67%	79%	63%
Viral Load Monitoring (2 Tests)	73%▲	58%	60%▲	47%	60%	47%
Syphilis Screening Rate	65%▲	58%	62%▲	54%	65%	52%
Controlling High Blood Pressure (Blood Pressure Reading < 140/90)	74%▲	67%	Rotated Off	N/A	69%	65%
Use of Spirometry Testing in the Assessment & Diagnosis of COPD	51%	46%	46%	45%	37%	42%
<i>Pharmacotherapy Management of COPD Exacerbation</i>						
Systemic Corticosteroid	54%▼	66%	61%	61%	68%	65%
Bronchodilator	82%	85%	82%	84%	85%	84%
Disease Modifying Anti-Rheumatic Drug Therapy in Rheumatoid Arthritis	76%	76%	73%	74%	74%	73%
Use of Imaging Studies for Low Back Pain	80%	79%	78%▼	80%	79%	N/A
<i>Annual Monitoring for Patients on Persistent Medications</i>						
ACE/ARBs	89%▼	91%	89%	90%	89%▲	88%
Digoxin	94%	94%	94%	91%	92%	91%
Diuretics	87%▼	90%	88%	88%	88%	87%
Anticonvulsants	62%▼	67%	61%▼	66%	66%	65%
Combined	87%▼	89%	88%	88%	88%▲	87%
<i>Antidepressant Medication Management</i>						
Effective Acute Treatment (84 days)	48%	52%	52%	51%	46%	49%
Effective Continuation Phase (180 days)	31%▼	35%	37%	35%	29%	32%
<i>Follow-up After Hospitalization for Mental Illness</i>						
Within 7 days	82%▲	70%	66%	68%	51%▼	63%
Within 30 days	91%▲	84%	80%	80%	72%▼	78%
<i>Follow-up Care for Children ADHD</i>						
Initiation Phase/ 30 Days	67%▲	58%	65%▲	56%	66%▲	54%
Continuation Phase/ 180 Days	82%▲	64%	79%▲	62%	82%▲	61%
<b>Acute Condition Care</b>						
Appropriate Treatment for Children w/ URI (Ages 3 Months to 18 Years)	92%▲	91%	92%▲	90%	91%▲	89%
Appropriate Testing for Children w/ Pharyngitis (Ages 2 to 18)	75%▼	84%	77%▼	83%	77%▼	81%
Avoidance of Antibiotic Treatment - Adults w/ Acute Bronchitis	26%	27%	22%▼	26%	23%	25%